



International Journal of Biology and Medicine

Research Article

Open Access

Hematological Parameters and Blood Smear effects of Tramadol on Male Rabbits

Fayrouz A khaled^{1*}, AMAAL A Younus² and Rui M Sale³

¹Chemistry Department, Faculty of Science, Omar Al-Mokhtar University, El -Beida-Libya

²Judicial experience& research center, El -Beida-Libya

³Faculty of Dentistry, Omar Al-Mokhtar University, El -Beida-Libya

*Corresponding Author: Fayrouz A Khaled, Chemistry Department, Faculty of Science, Omar Al-Mokhtar University, El -Beida-Libya, Email: faylzobair@yahoo.com

Received Date: Aug 04, 2022 / Accepted Date: Aug 20, 2022 / Published Date: Aug 25, 2022

Abstract

Analgesics are a bunch of drugs that act totally different ways on the central apprehensive framework to absence of pain and alleviation from torment. It is in this manner imperative to ponder the impacts of organization these analgesics to decide the side impacts for them utilize. Hence, we decided to study the hematological effects of tramadol on male rabbits. Tenth rabbits were arbitrarily partitioned into two breaks even with bunches (each bunch five rabbits). The primary gather was utilized as a control. The moment gather was utilized to consider the impact of tramadol (50 mg/kg body weight) for six weeks. Hemoglobin (Hb) substance, ruddy blood cells (RBCs) number and hematocrit (Hct) esteem diminished, whereas white blood cells (WBCs) and platelets tallies expanded after tramadol organization. Verbal organization of tramadol induced significant changes within the morphology of erythrocytes. In rabbit treated with tramadol the cells were appeared change in typical shape from star molded to sickle cells. The ponder appeared the destructive impacts of tramadol on the hematological parameters and blood spread on mail rabbits.

Keywords: Hematological Parameters; Tramadol; Rabbits

Cite this article as: Fayrouz A Khaled, AMAAL A Younus, Rui M Sale. 2022. Hematological Parameters and Blood Smear effects of Tramadol on Male Rabbits. Int J Biol Med. 4: 25-29.

Copyright: This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited. Copyright © 2022; Fayrouz A Khaled

Introduction

Opioids are the foremost capable painkillers. Their utilize is fitting for the treatment of moderate-to-severe persistent torment (5-10-point torment concentrated) with the objective to control this indication and to progress capacities and quality of life [1]. Tramadol may be a synthetic, centrally acting pain relieving, accessible in Europe since 1977 and within the Joined together States since 1995 for the treatment of torment disorders already amiable

as it were to the sedative analogues [2]. Tramadol has been in clinical utilize for the alleviation of mellow to direct torment in human and veterinary medication [3]. Tramadol is additionally utilized perioperatively in veterinary anesthesia because it essentially decreases the prerequisites of unstable anesthetics and opioid agents [4]. Tramadol, a engineered racemic blend of the 4-phenyl-piperidine simple of codeine, has gotten broad acknowledgment in human medication since it was to begin with presented in 1977 in

Germany [5]. These days, tramadol is broadly utilized as a pain relieving medicate in human medicine [6]. Its component of activity is based on the hindrance of climbing torment to the central anxious framework by its official to α -sedative receptors and hinder the reuptake of norepinephrine and serotonin [7]. Ceaseless tramadol organization leads to the appearance of its poisonous impacts on different organs of the body [8].

Materials and Method

Tested compounds

In this study tramadol was used. tramadol was purchased from pharmacy alsalam hospital in El -Beida-Libya.

Animals and treatments

Develop male Unused Zealand White rabbits age of 6 -week and introductory weight of (1892? 50.79 Kg) were utilized. Creatures were exclusively housed in cages and weighed week by week all through 6-week test period. Feed and water were provided ad libitum. Rabbits fed pellets which consisted of 30 % berseem (*Trifolium alexandrinum*) hay, 25 % yellow corn, 26.2% wheat bran, 14 % soybean meal, 3 % molasses, 1 % CaCl₂, 0.4 % NaCl, 0.3 % mixture of minerals and vitamins, and 0.1 % methionine. The vitamin and mineral premix per kg contained the following IU/gm for vitamins or minerals: vit A-4000,000, vit D3-5000, 000, vit E-16,7 g, K-0.67 g, vit B1-0.67 g, vit B2-2 g, B6-0.67 g, B12-0.004 g, B5-16.7 g, Pantothenic acid-6.67 g, Biotein-0.07 g, Folic acid-1.67 g, Choline chloride-400 g, Zn-23.3 g, Mn-10 g, Fe-25 g, Cu-1.67 g, I-0.25 g, Se-0.033 g, and Mg-133.4 g (Rabbit premix produced by Holland Feed Inter. Co.). The chemical analysis of the pellets[9] showed that they contained 15.8 % crude protein, 11.3 % crude fibre, 3.7 % ether extract, 7.2 % ash, 92.9 % organic matter and 62.4 % nitrogen free extract % as DM basis. The first group was used as control, while, groups 2 was treated with tramadol by gavage at a dose of 50 mg/kg

B.W/day (1/50 of tramadol) lethal dose[10]. for 6 successive weeks.

Hematological Parameters

Blood tests were collected from the ear vein of all creatures each week all through the 6-week exploratory period. Blood tests were gotten within the morning some time recently gets to to bolster and water. Values determined from total blood check (CBC). All CBC tests were performed by programmed blood cell analyzer (XP-300 Mechanized Hematology Analyzer, Sysmex American, Inc [11]. CBC were performed on EDTA as anti-coagulated tests. Differential cell tallies were performed physically utilizing Dif-Quik-stained blood smears. Information was recorded agreeing to the taking after categories: white blood cell (WBC); ruddy blood cell (RBC); hemoglobin (HB); cruel corpuscular volume (MCV); mean corpuscular hemoglobin (MCH) and mean corpuscular hemoglobin concentration (MCHC).

Preparation of blood smears

Lean movies, 3-5 cm in length, of the suctioned a drop of blood were made employing a smooth-edged glass spreader of not more than 2 cm in width. The blood parts are dragged behind the spreader and take off a path of cells behind them. After drying, films were dried within the discuss, and after that settled by drenching in a jolt of methanol for 15-20 min. Slides were exchanged to a recoloring bump containing Giemsa's recolor (1 volume of recolor with two volumes of refined water). After recoloring for 15-30 min, the slides were exchanged to a jostle containing buffered water, pH 7, quickly washed in three or four changes of water and at long last permitted to standing undisturbed in water for a brief time (as a rule 2-5 min) for separation to require put. Slides were exchanged to a recoloring jolt containing Giemsa's recolor (1 volume of recolor with two volumes of refined water). After recoloring for 15-30 min, the slides were exchanged to a jolt containing buffered water, pH 7, quickly

washed in three or four changes of water and at long last permitted to standing undisturbed in water for a brief time (more often than not 2-5 min) for separation to require put.

Result

Effect of tramadol on morphological pattern observed in blood smear

Oral administration of tramadol induced profound alterations in the morphology of erythrocytes. In rabbit treated with tramadol the cells were showed alteration in normal shape from star shaped to sickle cells. (Figure 1).




Figure 1: Photomicrographs showing peripheral blood smears of rabbits in two groups

Effect of tramadol on hematological parameters.

Table 1 represent the hematological parameters of male rabbits treated with tramadol (T). Results indicated that treatment with tramadol

significant decrease in red blood cells (RBC), white blood cells (WBC), packed cell volume (PCV), platelet count (PLT), hemoglobin (Hb), mean cell volume (MCV), mean cell hemoglobin (MCH) and mean cell hemoglobin concentration (MCHC).

Discussion

The show ponders appeared that tramadol caused diminish RBCS, Hb, PCV and PLT (Tables 1) assertion with [7], who found that organization of sildenafil (1.40 mg/kg/day) and

tramadol (4 mg/kg/day) for 25 days to rabbits driven to significant diminishes in RBCs tally and Hb substance. The diminish within the RBCs tally may be due to the inhibitory impact of tramadol on erythropoiesis [13]. In expansion, the WBCs check expanded after organization of tramadol to rats. [13] clarified the height within the WBCs tally by the enactment of defence resistant framework. Whereas [7] detailed that tall lymphocyte rate in male rats infused by tramadol (100 mg/kg) for 30 days was the reason behind the WBCs count elevation.

The ruddy blood cells (RBCs) tally appeared a common diminish in reaction to tramadol organization. This finding may be clarified on the premise of inhibitory impact of tramadol on histogenesis. The diminished in RBC number and haemoglobin (Hb) brought down the oxygen supply to distinctive tissues hence coming about in moo vitality generation.

Diminish in Hb contained MCH can be clarified due to diminished measure of RBC or impeded biosynthesis of home in bone marrow. These discoveries are in assertion with the detailed diminish in RBC check and Hb substance after treatment with tramadol [14].

Table 1: Changes Complete blood counts red blood cells RBCs, white blood cells WBCs, haemoglobin Hb, packed cell volume PCV, platelets, haemoglobin (Hb), and mean cell hemoglobin concentration (MCHC) and mean cell hemoglobin concentration (MCHC) of male rabbits treated with tramadol.

Parameter	Experimental groups	
	Control	Tramadol
RBC $\times 106 (\mu l)$	5.62 \pm 0.138 ^a	4.68 \pm 0.158 ^a
WBC $\times 103 (\mu l)$	8.53 \pm 0.18 ^a	7.41 \pm 0.35 ^b
HCT $\times 103 (\mu l)$	39.95 \pm 0.280 ^a	35.68 \pm 0.760 ^b
PLAT $\times 103 (\mu l)$	277.82 \pm 8.740 ^a	243.05 \pm 15.027 ^b
Hb (g/dl)	13.60 \pm 0.16 ^a	11.97 \pm 0.28 ^b
MCV (fl)	76.88 \pm 2.26 ^a	77.98 \pm 1.96 ^a
MCH (pg)	25.02 \pm 0.84 ^a	28.98 \pm 0.63 ^b
MCHC (dl)	32.69 \pm 0.45 ^a	32.82 \pm 0.57 ^b

Values are expressed as means \pm SE; n = 5 for each treatment group. Mean values within a row not sharing common superscript letters (a, b, c) were significantly different <0.05.

Conclusion

The comes about of the display ponder convincingly illustrated that tramadol presentation come about in shifting degree of haematological parameters of rabbits. Tramadol harmful impacts ought to be maintained a strategic distance from amid long term treatment extraordinarily in huge measurements.

Reference

- Huang AR., Mallet L. 2013. Prescribing opioids in older people. *Maturitas*. 74: 123-129. Ref.: <https://pubmed.ncbi.nlm.nih.gov/23201325/> Doi: <https://doi.org/10.1016/j.maturitas.2012.11.002>
- Moore KA, Cina SJ, Jones R, et al. 1999. Tissue distribution of tramadol and metabolites in an overdose fatality. *The American journal of forensic medicine and pathology*. 20: 98-100. Ref.: <https://pubmed.ncbi.nlm.nih.gov/10208348/> Doi: <https://doi.org/10.1097/00000433-199903000-00023>
- Pypendop BH, Ilkiw JE. 2008. Pharmacokinetics of tramadol, and its metabolite O-desmethyl tramadol, in cats. *Journal of veterinary pharmacology and therapeutics*. 31: 52-59. Ref.: <https://pubmed.ncbi.nlm.nih.gov/18177319/> Doi: <https://doi.org/10.1111/j.1365-2885.2007.00921.x>
- Seddighi MR., Egger CM, Rohrbach BW, et al. 2009. Effects of tramadol on the minimum alveolar concentration of sevoflurane in dogs. *Veterinary anaesthesia and analgesia*. 36: 334-340. Ref.: <https://pubmed.ncbi.nlm.nih.gov/19538570/>
- Li S, Perry CM. 2000. Tramadol: a review of its use in perioperative pain. *Drugs*. 60: 139-176. Ref.: <https://pubmed.ncbi.nlm.nih.gov/10929933/> Doi: <https://pubmed.ncbi.nlm.nih.gov/10929933/>

6. Azari O, Emadi L, Kheirandish R, et al. 2014. The effects of long-term administration of tramadol on epididymal sperm quality and testicular tissue in mice. *Iranian Journal of Veterinary Surgery*. 9: 23-30.?
7. Aldalou AR, Abdel-Aziz I, Shahwan O. 2014. Impact of giving sildenafil (viagra)/tramadol (tramal) combination on the blood of domestic rabbits. *Journal of Science*. 4: 162-169.?
8. Shadnia S, Soltaninejad K, Heydari K, et al. 2008. Tramadol intoxication: a review of 114 cases. *Human & experimental toxicology*. 27: 201-205. Ref.: <https://pubmed.ncbi.nlm.nih.gov/18650251/> Doi: <https://doi.org/10.1177/0960327108090270>
9. Smith JT, Mayer DT. 1955. Evaluation of sperm concentration by the hemacytometer method: Comparison of four counting fluids. *Fertility and Sterility*. 6: 271-275. Ref.: <https://pubmed.ncbi.nlm.nih.gov/14380385/> Doi: [https://doi.org/10.1016/s0015-0282\(16\)31987-2](https://doi.org/10.1016/s0015-0282(16)31987-2)
10. Ahmad RM, AL-Hubaity AY, Alazow NS. 2019. The Role of Vitamin C on the Structural Changes of Male Albino Rats Kidney Induced by Tramadol. *Annals of the College of Medicine, Mosul*. 41: 57-62.?
11. Turgeon M. 2012. *Clinical Hematology Theory and Procedures, Fundamentals of Hematological Analysis*. 5th ed, Philadelphia Lippincott Williams & Wilkins. 456-580.
12. Bain BJ. 2012. Blood cell morphology in health and disease. In *Dacie and Lewis Practical Haematology*. 69-100.
13. Aldiwan, MA, Alzobidy AMH, Younis MA. 2015. The effect of Tramadol on some blood and biochemical parameters of male rats (*Rattus norvegicus*). *Baghdad Science Journal*. 12.?
14. Goeringer KE, Logan BK, Christian GD. 1997. Identification of tramadol and its metabolites in blood from drug-related deaths and drug-impaired drivers. *Journal of analytical toxicology*. 21: 529-537. Ref.: <https://pubmed.ncbi.nlm.nih.gov/9399121/> Doi: <https://doi.org/10.1093/jat/21.7.529>